## JORDAN TYPE OF AN ARTINIAN ALGEBRA PROBLEM SESSIONS

## NANCY ABDALLAH AND PEDRO MACIAS MARQUES

## 1. Preliminaries

**Exercise 1.1.** Consider the Artinian algebra  $A = k[x, y]/(x^2, xy^2, y^5)$ , let m = (x, y) be its maximal ideal.

- (1) Find the Jordan types of  $y^2$ , xy, and  $xy + y^3$  in A.
- (2) Write down a Jordan basis for  $y^2$ , and the corresponding Jordan canonical form.
- (3) Describe the locus of the Jordan type (5, 2), i.e. the set

$$L_{(5,2)} = \{\ell \in \mathsf{m} : P_{\ell,A} = (5,2)\}.$$

(4) Suppose k is infinite. Which is the generic Jordan type of *A*?

**Exercise 1.2.** Consider the Artinian algebra  $A = k[x, y, z]/(yz - x^3, y^3, z^2)$ , a non-graded complete intersection.

- (1) Suppose char k = 2 and compute the Jordan type of  $\ell_1 = x + y + z$ .
- (2) Suppose chark  $\notin \{2,3,5\}$  and write  $\ell = ax + by + cz + \ell'$ , with  $\ell' \in m^2$ . Show that  $P_{\ell,A} = (7,5,3,3)$  if and only if  $ab \neq 0$ .

**Exercise 1.3.** Let *A* be an Artinian local algebra, with socle degree *j*, let m be its maximal ideal, and suppose that k = A/m is an infinite field.

- (1) Show that there is an element  $\ell \in m$  such that  $\ell^j \neq 0$ .
- (2) Show that if  $\ell' \in m^2$  then its Jordan type is not the generic Jordan type of *A*.

**Exercise 1.4.** Consider the algebra  $A = k[x, y]/(x^4, y^3)$ . Use the Lemma 2.11 to compute the Jordan type of x.

**Exercise 1.5.** Consider the algebra  $A = k[x, y, z]/(xy, xz, yz, x^j - y^j, x^j - z^j)$ . Compute the Jordan type of x.

**Exercise 1.6.** Let A = k[x, y]/I be an Artinian algebra, and let  $P_{x,A} = (p_1, \dots, p_s)$ . Consider the algebra B = k[x, y, z]/J, where  $J = (xz, yz, z^j, I)$ . What is the Jordan type  $P_{x,B}$ ?

## 2. Lefschetz properties

**Exercise 2.1.** Consider the Artinian algebra  $A = k[x, y]/(x^2, y^2)$ .

- (1) List all partitions of 4 that are dominated by  $H(A)^{\vee}$ .
- (2) Confirm that all partitions are attained as the Jordan type of an element in the maximal ideal.

**Exercise 2.2.** Consider the Artinian algebra  $A = k[x, y]/(x^2, xy^2, y^5)$ , let m = (x, y) be its maximal ideal. Give an example of a partition that is dominated by  $H(A)^{\vee}$  but is not the Jordan type of any element  $\ell \in m$ .

3. Artinian Gorenstein Algebras

Exercise 3.1. Consider the Artinian Gorenstein algebra

$$A = k\{x, y\} / (x^3 - y^5, x^2 y^2).$$

Write down a monomial basis for A, where the order of each element is apparent (i.e if  $m_1, m_2 \in k\{x, y\}$  are two monomials such that  $m_1 = m_2$  in A, choose the one with highest degree).

**Exercise 3.2.** Consider the Artinian Gorenstein algebra  $A = k\{x, y, z\} / \operatorname{Ann} F$ , for  $F = X^2 Y^2 + Z^3$ .

- (1) Compute its Q(a) modules, and get the Hilbert function of A as the sum of its symmetric decomposition.
- (2) Check that Ann  $F = (xz, yz, x^3, y^3, z^3 x^2y^2)$ .
- (3) Compute the Jordan type  $P_{x+y+z,A}$ . Is this a weak or strong Lefschetz element?
- (4) Find two Jordan types of elements of order one (i.e. elements in  $m \setminus m^2$ ) that are not comparable.

Exercise 3.3. Let *A* be an Artinian Gorenstein algebra, with maximal ideal m.

$$z \in \mathsf{m}^k \cap (0 : \mathsf{m}^c)$$
, and  $\ell \in \mathsf{m}^i$ , with  $c > i$ . Show that

$$\ell z \in \mathsf{m}^{i+k} \cap (0:\mathsf{m}^{c-i}).$$

(2) Suppose *z* represents a non-zero class in  $Q(a)_k$  and  $\ell \in \mathsf{m}^i$ . Show that if  $\ell z$  represents a non-zero class in  $Q(b)_m$ , and b < a then  $m \ge k + 2$ .

**Exercise 3.4.** Let *A* be an Artinian Gorenstein algebra with Hilbert function

$$H(A) = (1, 3, 3, 4, 2, 1)$$

- (1) Compute the Hilbert functions of its Q(a) modules.
- (2) Use Exercise 3.3 to show that *A* is not strong Lefschetz.

(1) Let