

JORDAN TYPE OF AN ARTINIAN ALGEBRA PROBLEM SESSIONS

NANCY ABDALLAH AND PEDRO MACIAS MARQUES

1. PRELIMINARIES

Exercise 1.1. Consider the Artinian algebra $A = \mathbb{k}[x, y]/(x^2, xy^2, y^5)$, let $\mathfrak{m} = (x, y)$ be its maximal ideal.

- (1) Find the Jordan types of y^2 , xy , and $xy + y^3$ in A .
- (2) Write down a Jordan basis for y^2 , and the corresponding Jordan canonical form.
- (3) Describe the locus of the Jordan type $(5, 2)$, i.e. the set

$$L_{(5,2)} = \{\ell \in \mathfrak{m} : P_{\ell,A} = (5, 2)\}.$$

- (4) Suppose \mathbb{k} is infinite. Which is the generic Jordan type of A ?

Exercise 1.2. Consider the Artinian algebra $A = \mathbb{k}[x, y, z]/(yz - x^3, y^3, z^2)$, a non-graded complete intersection.

- (1) Suppose $\text{char } \mathbb{k} = 2$ and compute the Jordan type of $\ell_1 = x + y + z$.
- (2) Suppose $\text{char } \mathbb{k} \notin \{2, 3, 5\}$ and write $\ell = ax + by + cz + \ell'$, with $\ell' \in \mathfrak{m}^2$. Show that $P_{\ell,A} = (7, 5, 3, 3)$ if and only if $ab \neq 0$.

Exercise 1.3. Let A be an Artinian local algebra, with socle degree j , let \mathfrak{m} be its maximal ideal, and suppose that $\mathbb{k} = A/\mathfrak{m}$ is an infinite field.

- (1) Show that there is an element $\ell \in \mathfrak{m}$ such that $\ell^j \neq 0$.
- (2) Show that if $\ell' \in \mathfrak{m}^2$ then its Jordan type is not the generic Jordan type of A .

Exercise 1.4. Consider the algebra $A = \mathbb{k}[x, y]/(x^4, y^3)$. Use the Lemma 2.11 to compute the Jordan type of x .

Exercise 1.5. Consider the algebra $A = \mathbb{k}[x, y, z]/(xy, xz, yz, x^j - y^j, x^j - z^j)$. Compute the Jordan type of x .

Exercise 1.6. Let $A = \mathbb{k}[x, y]/I$ be an Artinian algebra, and let $P_{x,A} = (p_1, \dots, p_s)$. Consider the algebra $B = \mathbb{k}[x, y, z]/J$, where $J = (xz, yz, z^j, I)$. What is the Jordan type $P_{x,B}$?

2. LEFSCHETZ PROPERTIES

Exercise 2.1. Consider the Artinian algebra $A = \mathbb{k}[x, y]/(x^2, y^2)$.

- (1) List all partitions of 4 that are dominated by $H(A)^\vee$.
- (2) Confirm that all partitions are attained as the Jordan type of an element in the maximal ideal.

Exercise 2.2. Consider the Artinian algebra $A = k[x, y]/(x^2, xy^2, y^5)$, let $\mathfrak{m} = (x, y)$ be its maximal ideal. Give an example of a partition that is dominated by $H(A)^\vee$ but is not the Jordan type of any element $\ell \in \mathfrak{m}$.